

PATENT

Attorney Docket No. 6794S-000010/US

Client Docket No. K-00367

Express Mail Label No. EL 838754002 US

APPLICATION FOR LETTERS PATENT

for

**ENHANCED PHARMACOKINETIC PROFILE OF INTRADERMALLY
DELIVERED SUBSTANCES**

09897304.052904
T05250 "T032500

1997-1998 1998-1999 1999-2000 2000-2001 2001-2002 2002-2003 2003-2004 2004-2005 2005-2006 2006-2007 2007-2008 2008-2009 2009-2010 2010-2011 2011-2012 2012-2013 2013-2014 2014-2015 2015-2016 2016-2017 2017-2018 2018-2019 2019-2020 2020-2021 2021-2022 2022-2023 2023-2024 2024-2025 2025-2026 2026-2027 2027-2028 2028-2029 2029-2030 2030-2031 2031-2032 2032-2033 2033-2034 2034-2035 2035-2036 2036-2037 2037-2038 2038-2039 2039-2040 2040-2041 2041-2042 2042-2043 2043-2044 2044-2045 2045-2046 2046-2047 2047-2048 2048-2049 2049-2050 2050-2051 2051-2052 2052-2053 2053-2054 2054-2055 2055-2056 2056-2057 2057-2058 2058-2059 2059-2060 2060-2061 2061-2062 2062-2063 2063-2064 2064-2065 2065-2066 2066-2067 2067-2068 2068-2069 2069-2070 2070-2071 2071-2072 2072-2073 2073-2074 2074-2075 2075-2076 2076-2077 2077-2078 2078-2079 2079-2080 2080-2081 2081-2082 2082-2083 2083-2084 2084-2085 2085-2086 2086-2087 2087-2088 2088-2089 2089-2090 2090-2091 2091-2092 2092-2093 2093-2094 2094-2095 2095-2096 2096-2097 2097-2098 2098-2099 2099-2100 2100-2101 2101-2102 2102-2103 2103-2104 2104-2105 2105-2106 2106-2107 2107-2108 2108-2109 2109-2110 2110-2111 2111-2112 2112-2113 2113-2114 2114-2115 2115-2116 2116-2117 2117-2118 2118-2119 2119-2120 2120-2121 2121-2122 2122-2123 2123-2124 2124-2125 2125-2126 2126-2127 2127-2128 2128-2129 2129-2130 2130-2131 2131-2132 2132-2133 2133-2134 2134-2135 2135-2136 2136-2137 2137-2138 2138-2139 2139-2140 2140-2141 2141-2142 2142-2143 2143-2144 2144-2145 2145-2146 2146-2147 2147-2148 2148-2149 2149-2150 2150-2151 2151-2152 2152-2153 2153-2154 2154-2155 2155-2156 2156-2157 2157-2158 2158-2159 2159-2160 2160-2161 2161-2162 2162-2163 2163-2164 2164-2165 2165-2166 2166-2167 2167-2168 2168-2169 2169-2170 2170-2171 2171-2172 2172-2173 2173-2174 2174-2175 2175-2176 2176-2177 2177-2178 2178-2179 2179-2180 2180-2181 2181-2182 2182-2183 2183-2184 2184-2185 2185-2186 2186-2187 2187-2188 2188-2189 2189-2190 2190-2191 2191-2192 2192-2193 2193-2194 2194-2195 2195-2196 2196-2197 2197-2198 2198-2199 2199-2200 2200-2201 2201-2202 2202-2203 2203-2204 2204-2205 2205-2206 2206-2207 2207-2208 2208-2209 2209-2210 2210-2211 2211-2212 2212-2213 2213-2214 2214-2215 2215-2216 2216-2217 2217-2218 2218-2219 2219-2220 2220-2221 2221-2222 2222-2223 2223-2224 2224-2225 2225-2226 2226-2227 2227-2228 2228-2229 2229-2230 2230-2231 2231-2232 2232-2233 2233-2234 2234-2235 2235-2236 2236-2237 2237-2238 2238-2239 2239-2240 2240-2241 2241-2242 2242-2243 2243-2244 2244-2245 2245-2246 2246-2247 2247-2248 2248-2249 2249-2250 2250-2251 2251-2252 2252-2253 2253-2254 2254-2255 2255-2256 2256-2257 2257-2258 2258-2259 2259-2260 2260-2261 2261-2262 2262-2263 2263-2264 2264-2265 2265-2266 2266-2267 2267-2268 2268-2269 2269-2270 2270-2271 2271-2272 2272-2273 2273-2274 2274-2275 2275-2276 2276-2277 2277-2278 2278-2279 2279-2280 2280-2281 2281-2282 2282-2283 2283-2284 2284-2285 2285-2286 2286-2287 2287-2288 22	
---	--

[0001] The present invention relates to methods and devices for administration of substances into the intradermal layer of skin.

BACKGROUND OF THE INVENTION

[0002] The importance of efficiently and safely administering pharmaceutical substances such as diagnostic agents and drugs has long been recognized. Although an important consideration for all pharmaceutical substances, obtaining adequate bioavailability of large molecules such as proteins that have arisen out of the biotechnology industry has recently highlighted this need to obtain efficient and reproducible absorption (Cleland et al., *Curr. Opin. Biotechnol.* 12: 212-219, 2001). The use of conventional needles has long provided one approach for delivering pharmaceutical substances to humans and animals by administration through the skin. Considerable effort has been made to achieve reproducible and efficacious delivery through the skin while improving the ease of injection and reducing patient apprehension and/or pain associated with conventional needles. Furthermore, certain delivery systems eliminate needles entirely, and rely upon chemical mediators or external driving forces such as iontophoretic currents or electroporation or thermal poration or sonophoresis to breach the stratum corneum, the outermost layer of the skin, and deliver substances through the surface of the skin. However, such delivery systems do not reproducibly breach the skin barriers or deliver the pharmaceutical substance to a given depth below the surface of the skin and consequently, clinical results can be variable. Thus, mechanical breach of the stratum corneum such as with needles, is believed to provide the most reproducible method of administration of substances through the surface of the skin, and to provide control and reliability in placement of administered substances.

[0003] Approaches for delivering substances beneath the surface of the skin have almost exclusively involved transdermal administration, i.e. delivery of substances through the skin to a site beneath the skin. Transdermal delivery includes subcutaneous, intramuscular or intravenous routes of administration of which,

intramuscular (IM) and subcutaneous (SC) injections have been the most commonly used .

[0004] Anatomically, the outer surface of the body is made up of two major tissue layers, an outer epidermis and an underlying dermis, which together constitute the skin (for review, see *Physiology, Biochemistry, and Molecular Biology of the Skin, Second Edition*, L.A. Goldsmith, Ed., Oxford University Press, New York, 1991). The epidermis is subdivided into five layers or strata of a total thickness of between 75 and 150 μm . Beneath the epidermis lies the dermis, which contains two layers, an outermost portion referred to as the papillary dermis and a deeper layer referred to as the reticular dermis. The papillary dermis contains vast microcirculatory blood and lymphatic plexuses. In contrast, the reticular dermis is relatively acellular and avascular and made up of dense collagenous and elastic connective tissue. Beneath the epidermis and dermis is the subcutaneous tissue, also referred to as the hypodermis, which is composed of connective tissue and fatty tissue. Muscle tissue lies beneath the subcutaneous tissue.

[0005] As noted above, both the subcutaneous tissue and muscle tissue have been commonly used as sites for administration of pharmaceutical substances. The dermis, however, has rarely been targeted as a site for administration of substances, and this may be due, at least in part, to the difficulty of precise needle placement into the intradermal space. Furthermore, even though the dermis, in particular, the papillary dermis has been known to have a high degree of vascularity, it has not heretofore been appreciated that one could take advantage of this high degree of vascularity to obtain an improved absorption profile for administered substances compared to subcutaneous administration. This is because small drug molecules are typically rapidly absorbed after administration into the subcutaneous tissue which has been far more easily and predictably targeted than the dermis has been. On the other hand, large molecules such as proteins are typically not well absorbed through the capillary epithelium regardless of the degree of vascularity so that one would not have expected to achieve a significant absorption advantage over subcutaneous

administration by the more difficult to achieve intradermal administration even for large molecules.

[0006] One approach to administration beneath the surface to the skin and into the region of the intradermal space has been routinely used in the Mantoux tuberculin test. In this procedure, a purified protein derivative is injected at a shallow angle to the skin surface using a 27 or 30 gauge needle (Flynn et al, *Chest* 106: 1463-5, 1994). A degree of uncertainty in placement of the injection can, however, result in some false negative test results. Moreover, the test has involved a localized injection to elicit a response at the site of injection and the Mantoux approach has not led to the use of intradermal injection for systemic administration of substances.

[0007] Some groups have reported on systemic administration by what has been characterized as "intradermal" injection. In one such report, a comparison study of subcutaneous and what was described as "intradermal" injection was performed (Autret et al, *Therapie* 46:5-8, 1991). The pharmaceutical substance tested was calcitonin, a protein of a molecular weight of about 3600. Although it was stated that the drug was injected intradermally, the injections used a 4 mm needle pushed up to the base at an angle of 60. This would have resulted in placement of the injectate at a depth of about 3.5 mm and into the lower portion of the reticular dermis or into the subcutaneous tissue rather than into the vascularized papillary dermis. If, in fact, this group injected into the lower portion of the reticular dermis rather than into the subcutaneous tissue, it would be expected that the substance would either be slowly absorbed in the relatively less vascular reticular dermis or diffuse into the subcutaneous region to result in what would be functionally the same as subcutaneous administration and absorption. Such actual or functional subcutaneous administration would explain the reported lack of difference between subcutaneous and what was characterized as intradermal administration, in the times at which maximum plasma concentration was reached, the concentrations at each assay time and the areas under the curves.

[0008] Similarly, Bressolle et al. administered sodium ceftazidime in what was characterized as "intra-dermal" injection using a 4 mm needle (Bressolle et al. *J. Pharm. Sci.* 82:1175-1178, 1993). This would have resulted in injection to a depth of 4 mm below the skin surface to produce actual or functional subcutaneous injection, although good subcutaneous absorption would have been anticipated in this instance because sodium ceftazidime is hydrophilic and of relatively low molecular weight.

[0009] Another group reported on what was described as an intra-dermal drug delivery device (U.S. Patent No. 5,007,501). Injection was indicated to be at a slow rate and the injection site was intended to be in some region below the epidermis, i.e., the interface between the epidermis and the dermis or the interior of the dermis or subcutaneous tissue. This reference, however, provided no teachings that would suggest a selective administration into the dermis nor did the reference suggest any possible pharmacokinetic advantage that might result from such selective administration.

[0010] Thus there remains a continuing need for efficient and safe methods and devices for administration of pharmaceutical substances.

SUMMARY OF THE INVENTION.

[0011] The present disclosure relates to a new parenteral administration method based on directly targeting the dermal space whereby such method dramatically alters the pharmacokinetics (PK) and pharmacodynamics (PD) parameters of administered substances. By the use of direct intra-dermal (ID) administration means hereafter referred to as dermal-access means, for example, using microneedle-based injection and infusion systems (or other means to accurately target the intra-dermal space), the pharmacokinetics of many substances including drugs and diagnostic substances, which are especially protein and peptide hormones, can be altered when compared to traditional parental administration routes of subcutaneous and intravenous delivery. These findings are pertinent not only to microdevice-based injection means, but other

delivery methods such as needleless or needle-free ballistic injection of fluids or powders into the ID space, Mantoux-type ID injection, enhanced iontophoresis through microdevices, and direct deposition of fluid, solids, or other dosing forms into the skin. Disclosed is a method to increase the rate of uptake for parenterally-administered drugs without necessitating IV access. One significant beneficial effect of this delivery method is providing a shorter T_{max} (time to achieve maximum blood concentration of the drug). Potential corollary benefits include higher maximum concentrations for a given unit dose (C_{max}), higher bioavailability, more rapid uptake rates, more rapid onset of pharmacodynamics or biological effects, and reduced drug depot effects. According to the present invention, improved pharmacokinetics means increased bioavailability, decreased lag time (T_{lag}), decreased T_{max} , more rapid absorption rates, more rapid onset and/or increased C_{max} for a given amount of compound administered, compared to subcutaneous, intramuscular or other non-IV parenteral means of drug delivery.

[0012] By bioavailability is meant the total amount of a given dosage that reached the blood compartment. This is generally measured as the area under the curve in a plot of concentration vs. time. By "lag time" is meant the delay between the administration of a compound and time to measurable or detectable blood or plasma levels. T_{max} is a value representing the time to achieve maximal blood concentration of the compound, and C_{max} is the maximum blood concentration reached with a given dose and administration method. The time for onset is a function of T_{lag} , T_{max} and C_{max} , as all of these parameters influence the time necessary to achieve a blood (or target tissue) concentration necessary to realize a biological effect. T_{max} and C_{max} can be determined by visual inspection of graphical results and can often provide sufficient information to compare two methods of administration of a compound. However, numerical values can be determined more precisely by analysis using kinetic models (as described below) and/or other means known to those of skill in the art.

under 2 to about 4 mm in other regions of the body depending upon the particular study report (Hwang et al., *Ann Plastic Surg* 46:327-331, 2001; Southwood, *Plast. Reconstr. Surg* 15:423-429, 1955; Rushmer et al., *Science* 154:343-348, 1966).

[0015] As used herein, intradermal is intended to mean administration of a substance into the dermis in such a manner that the substance readily reaches the richly vascularized papillary dermis and is rapidly absorbed into the blood capillaries and/or lymphatic vessels to become systemically bioavailable. Such can result from placement of the substance in the upper region of the dermis, i.e. the papillary dermis or in the upper portion of the relatively less vascular reticular dermis such that the substance readily diffuses into the papillary dermis. It is believed that placement of a substance predominately at a depth of at least about 0.3 mm, more preferably, at least about 0.4 mm and most preferably at least about 0.5 mm up to a depth of no more than about 2.5 mm, more preferably, no more than about 2.0 mm and most preferably no more than about 1.7 mm will result in rapid absorption of macromolecular and/or hydrophobic substances. Placement of the substance predominately at greater depths and/or into the lower portion of the reticular dermis is believed to result in the substance being slowly absorbed in the less vascular reticular dermis or in the subcutaneous region either of which would result in reduced absorption of macromolecular and/or hydrophobic substances. The controlled delivery of a substance in this dermal space below the papillary dermis in the reticular dermis, but sufficiently above the interface between the dermis and the subcutaneous tissue, should enable an efficient (outward) migration of the substance to the (undisturbed) vascular and lymphatic microcapillary bed (in the papillary dermis), where it can be absorbed into systemic circulation via these microcapillaries without being sequestered in transit by any other cutaneous tissue compartment.

[0016] Another benefit of the invention is to achieve more rapid systemic distribution and offset of drugs or diagnostic agents. This is also pertinent for many hormones that in the body are secreted in a pulsatile fashion. Many side effects are associated with having continuous circulating levels of substances administered. A

very pertinent example is female reproductive hormones that actually have the opposite effect (cause infertility) when continuously present in the blood. Likewise, continuous and elevated levels of insulin are suspected to down regulate insulin receptors both in quantity and sensitivity.

[0017] Another benefit of the invention is to achieve higher bioavailabilities of drugs or diagnostic agents. This effect has been most dramatic for ID administration of high molecular weight substances, especially proteins, peptides, and polysaccharides. The direct benefit is that ID administration with enhanced bioavailability, allows equivalent biological effects while using less active agent. This results in direct economic benefit to the drug manufacturer and perhaps consumer, especially for expensive protein therapeutics and diagnostics. Likewise, higher bioavailability may allow reduced overall dosing and decrease the patient's side effects associated with higher dosing.

[0018] Another benefit of the invention is the attainment of higher maximum concentrations of drugs or diagnostic substances. The inventors have found that substances administered ID are absorbed more rapidly, with bolus administration resulting in higher initial concentrations. This is most beneficial for substances whose efficacy is related to maximal concentration. The more rapid onset allows higher C_{Max} values to be reached with lesser amounts of the substance. Therefore, the dose can be reduced, providing an economic benefit, as well as a physiological benefit since lesser amounts of the drug or diagnostic agent has to be cleared by the body.

[0019] Another benefit of the invention is no change in systemic elimination rates or intrinsic clearance mechanisms of drugs or diagnostic agents. All studies to date by the applicants have maintained the same systemic elimination rate for the substances tested as via IV or SC dosing routes. This indicates this dosing route has no change in the biological mechanism for systemic clearance. This is an advantageous from a regulatory standpoint, since degradation and clearance pathways

need not be reinvestigated prior to filing for FDA approval. This is also beneficial from a pharmacokinetics standpoint, since it allows predictability of dosing regimes. Some substances may be eliminated from the body more rapidly if their clearance mechanism are concentration dependent. Since ID delivery results in higher C_{max} , clearance rate may be increased, although the intrinsic mechanism remains unchanged.

[0020] Another benefit of the invention is no change in pharmacodynamic mechanism or biological response mechanism. As stated above, administered drugs by the methods taught by the applicants still exert their effects by the same biological pathways that are intrinsic to other delivery means. Any pharmacodynamic changes are related only to the difference patterns of appearance, disappearance, and drug or diagnostic agent concentrations present in the biological system.

[0021] Using the methods of the present invention, pharmaceutical compounds may be administered as a bolus, or by infusion. As used herein, the term "bolus" is intended to mean an amount that is delivered within a time period of less than ten (10) minutes. "Infusion" is intended to mean the delivery of a substance over a time period greater than ten (10) minutes. It is understood that bolus administration or delivery can be carried out with rate controlling means, for example a pump, or have no specific rate controlling means, for example user self-injection.

[0022] Another benefit of the invention is removal of the physical or kinetic barriers invoked when drugs passes through and becomes trapped in cutaneous tissue compartments prior to systemic absorption. Elimination of such barriers leads to an extremely broad applicability to various drug classes. Many drugs administered subcutaneously exert this depot effect -- that is, the drug is slowly released from the SC space, in which it is trapped, as the rate determining step prior to systemic absorption, due to affinity for or slow diffusion through the fatty adipose tissue. This depot effect results in a lower C_{max} and longer T_{max} , compared to ID, and can result in high inter-individual variability of absorption. This effect is also pertinent for

particularly good for substances such as growth hormones, or analgesics. Longer duration infusion, typically at lower infusion rates can result in continuous low basal levels of drugs that is desired for anticoagulants, basal insulin, and chronic pain therapy. These kinetic profiles can be combined in multiple fashion to exhibit almost any kinetic profile desired. An example would be to pulsatile delivery of fertility hormone (LHRH) for pregnancy induction, which requires intermittent peaks every 90 minutes with total clearance between pulses. Other examples would be rapid peak onset of drugs for migraine relief, followed by lower levels for pain prophylaxis.

[0024] Another benefit of the invention is reduced degradation of drugs and diagnostic agents and/or undesirable immunogenic activity. Transdermal methods using chemical enhancers or iontophoresis, or sonophoresis or electroporation or thermal poration require that a drug pass through the viable epidermal layer, which has high metabolic and immunogenic activity. Metabolic conversion of substances in the epidermis or sequestration by immunoglobulins reduces the amount of drug available for absorption. The ID administration circumvents this problem by placing the drug directly in the dermis, thus bypassing the epidermis entirely.

[0025] These and other benefits of the invention are achieved by directly targeting absorption by the papillary dermis and by controlled delivery of drugs, diagnostic agents, and other substances to the dermal space of skin. The inventors have found that by specifically targeting the intradermal space and controlling the rate and pattern of delivery, the pharmacokinetics exhibited by specific drugs can be unexpectedly improved, and can in many situations be varied with resulting clinical advantage. Such pharmacokinetics cannot be as readily obtained or controlled by other parenteral administration routes, except by IV access.

[0026] The present invention improves the clinical utility of ID delivery of drugs, diagnostic agents, and other substances to humans or animals. The methods employ dermal-access means (for example a small gauge needle, especially microneedles), to directly target the intradermal space and to deliver substances to the intradermal space

as a bolus or by infusion. It has been discovered that the placement of the dermal-access means within the dermis provides for efficacious delivery and pharmacokinetic control of active substances. The dermal-access means is so designed as to prevent leakage of the substance from the skin and improve adsorption within the intradermal space. The pharmacokinetics of hormone drugs delivered according to the methods of the invention have been found to be vastly different to the pharmacokinetics of conventional SC delivery of the drug, indicating that ID administration according to the methods of the invention will provide improved clinical results. Delivery devices that place the dermal-access means at an appropriate depth in the intradermal space and control the volume and rate of fluid delivery provide accurate delivery of the substance to the desired location without leakage.

[0027] . Disclosed is a method to increase the rate of uptake for parenterally-administered drugs without necessitating IV access. This effect provides a shorter T_{max} . Potential corollary benefits include higher maximum concentrations for a given unit dose (C_{max}), higher bioavailability, more rapid onset of pharmacodynamics or biological effects, and reduced drug depot effects.

[0028] It has also been found that by appropriate depth control of the dermal-access means within the intradermal space that the pharmacokinetics of hormone drugs delivered according to the methods of the invention can, if required, produce similar clinical results to that of conventional SC delivery of the drug.

[0029] The pharmacokinetic profile for individual compounds will vary according to the chemical properties of the compounds. For example, compounds that are relatively large, having a molecular weight of at least 1000 Daltons as well as larger compounds of at least 2000 Daltons, at least 4000 Daltons, at least 10,000 Daltons and larger and/or hydrophobic compounds are expected to show the most significant changes compared to traditional parenteral methods of administration, such as intramuscular, subcutaneous or subdermal injection. It is expected that small

hydrophilic substances, on the whole, will exhibit similar kinetics for ID delivery compared to other methods.

DESCRIPTION OF THE DRAWINGS

[0030] Figure 1 shows a timecourse of plasma insulin levels of intradermal versus subcutaneous bolus administration of fast-acting.

[0031] Figure 2 shows a timecourse of blood glucose levels of intradermal versus subcutaneous bolus administration of fast-acting insulin.

[0032] Figure 3 shows a comparison of bolus ID dosing of fast-acting versus regular insulin.

[0033] Figure 4 shows the effects of different intradermal injection depths for bolus dosing of fast-acting insulin on the timecourse of insulin levels

[0034] Figure 5 shows a comparison of the timecourse of insulin levels for bolus dosing of long-acting insulin administered subcutaneously or intradermally.

[0035] Figure 6 and 7 show a comparison of the pharmacokinetic availability and the pharmacodynamic results of granulocyte colony stimulating factor delivered intradermally with a single needle or three point needle array, subcutaneously, or intravenously.

[0036] Figures 8, 9 and 10 show a comparison of low molecular weight heparin intradermal delivery by bolus, short duration, long duration infusion with comparison to subcutaneous infusion.

[0037] Figure 11 shows a timecourse of plasma genotropin levels of intradermal single needle, intradermal array and subcutaneous bolus administration.

DETAILED DESCRIPTION OF THE INVENTION

[0038] The present invention provides a method for therapeutic treatment by delivery of a drug or other substance to a human or animal subject by directly targeting the intradermal space, where the drug or substance is administered to the intradermal space through one or more dermal-access means incorporated within the device. Substances infused according to the methods of the invention have been

found to exhibit pharmacokinetics superior to, and more clinically desirable than that observed for the same substance administered by SC injection.

[0039] The dermal-access means used for ID administration according to the invention is not critical as long as it penetrates the skin of a subject to the desired targeted depth within the intradermal space without passing through it. In most cases, the device will penetrate the skin and to a depth of about 0.5-2 mm. The dermal-access means may comprise conventional injection needles, catheters or microneedles of all known types, employed singularly or in multiple needle arrays. The dermal-access means may comprise needleless devices including ballistic injection devices. The terms "needle" and "needles" as used herein are intended to encompass all such needle-like structures. The term microneedles as used herein are intended to encompass structures smaller than about 30 gauge, typically about 31-50 gauge when such structures are cylindrical in nature. Non-cylindrical structures encompass by the term microneedles would therefore be of comparable diameter and include pyramidal, rectangular, octagonal, wedged, and other geometrical shapes. Dermal-access means also include ballistic fluid injection devices, powder-jet delivery devices, piezoelectric, electromotive, electromagnetic assisted delivery devices, gas-assisted delivery devices, of which directly penetrate the skin to provide access for delivery or directly deliver substances to the targeted location within the dermal space. By varying the targeted depth of delivery of substances by the dermal-access means, pharmacokinetic and pharmacodynamic (PK/PD) behavior of the drug or substance can be tailored to the desired clinical application most appropriate for a particular patient's condition. The targeted depth of delivery of substances by the dermal-access means may be controlled manually by the practitioner, or with or without the assistance of indicator means to indicate when the desired depth is reached. Preferably however, the device has structural means for controlling skin penetration to the desired depth within the intradermal space. This is most typically accomplished by means of a widened area or hub associated with the shaft of the dermal-access means that may take the form of a backing structure or platform to which the needles are attached. The length of microneedles as dermal-access means

are easily varied during the fabrication process and are routinely produced in less than 2 mm length. Microneedles are also a very sharp and of a very small gauge, to further reduce pain and other sensation during the injection or infusion. They may be used in the invention as individual single-lumen microneedles or multiple microneedles may be assembled or fabricated in linear arrays or two-dimensional arrays as to increase the rate of delivery or the amount of substance delivered in a given period of time. Microneedles may be incorporated into a variety of devices such as holders and housings that may also serve to limit the depth of penetration. The dermal-access means of the invention may also incorporate reservoirs to contain the substance prior to delivery or pumps or other means for delivering the drug or other substance under pressure. Alternatively, the device housing the dermal-access means may be linked externally to such additional components.

[0040] IV-like pharmacokinetics is accomplished by administering drugs into the dermal compartment in intimate contact with the capillary microvasculature and lymphatic microvasculature. It should be understood that the terms microcapillaries or capillary beds refer to either vascular or lymphatic drainage pathways within the dermal area.

[0041] While not intending to be bound by any theoretical mechanism of action, it is believed that the rapid absorption observed upon administration into the dermis is achieved as a result of the rich plexuses of blood and lymphatic vessels in the dermis. However, the presence of blood and lymphatic plexuses in the dermis would not by itself be expected to produce an enhanced absorption of macromolecules. This is because capillary endothelium is normally of low permeability or impermeable to macromolecules such as proteins, polysaccharides, nucleic acid polymers, substance having polymers attached such as pegylated proteins and the like. Such macromolecules have a molecular weight of at least 1000 Daltons or of a higher molecular weight of at least, 2000 Daltons, at least 4000 Daltons, at least 10,000 Daltons or even higher. Furthermore, a relatively slow lymphatic drainage from the interstitium into the vascular compartment would also not be expected to produce a

rapid increase in plasma concentration upon placement of a pharmaceutical substance into the dermis.

[0042] One possible explanation for the unexpected enhanced absorption reported herein is that upon injection of substances so that they readily reach the papillary dermis an increase in blood flow and capillary permeability results. For example, it is known that a pinprick insertion to a depth of 3 mm produces an increase in blood flow and this has been postulated to be independent of pain stimulus and due to tissue release of histamine (Arildsson et al., *Microvascular Res.* 59:122-130, 2000). This is consistent with the observation that an acute inflammatory response elicited in response to skin injury produces a transient increase in blood flow and capillary permeability (see *Physiology, Biochemistry, and Molecular Biology of the Skin, Second Edition*, L.A. Goldsmith, Ed., Oxford Univ. Press, New York, 1991, p. 1060; Wilhem, *Rev. Can. Biol.* 30:153-172, 1971). At the same time, the injection into the intradermal layer would be expected to increase interstitial pressure. It is known that increasing interstitial pressure from values (beyond the "normal range") of about -7 to about +2 mmHg distends lymphatic vessels and increases lymph flow (Skobe et al., *J. Investig. Dermatol. Symp. Proc.* 5:14-19, 2000). Thus, the increased interstitial pressure elicited by injection into the intradermal layer is believed to elicit increased lymph flow and increased absorption of substances injected into the dermis.

[0043] By "improved pharmacokinetics" it is meant that an enhancement of pharmacokinetic profile is achieved as measured, for example, by standard pharmacokinetic parameters such as time to maximal plasma concentration (T_{max}), the magnitude of maximal plasma concentration (C_{max}) or the time to elicit a minimally detectable blood or plasma concentration (T_{lag}). By enhanced absorption profile, it is meant that absorption is improved or greater as measured by such pharmacokinetic parameters. The measurement of pharmacokinetic parameters and determination of minimally effective concentrations are routinely performed in the art. Values obtained are deemed to be enhanced by comparison with a standard route of administration such as, for example, subcutaneous administration or intramuscular

polymers, sugars, biological and or biodegradable materials, and/or combinations thereof.

[0046] It has been found that certain features of the intradermal administration methods provide clinically useful PK/PD and dose accuracy. For example, it has been found that placement of the needle outlet within the skin significantly affects PK/PD parameters. The outlet of a conventional or standard gauge needle with a bevel has a relatively large exposed height (the vertical rise of the outlet). Although the needle tip may be placed at the desired depth within the intradermal space, the large exposed height of the needle outlet causes the delivered substance to be deposited at a much shallower depth nearer to the skin surface. As a result, the substance tends to effuse out of the skin due to backpressure exerted by the skin itself and to pressure built up from accumulating fluid from the injection or infusion. That is, at a greater depth a needle outlet with a greater exposed height will still seal efficiently where as an outlet with the same exposed height will not seal efficiently when placed in a shallower depth within the intradermal space. Typically, the exposed height of the needle outlet will be from 0 to about 1 mm. A needle outlet with an exposed height of 0 mm has no bevel and is at the tip of the needle. In this case, the depth of the outlet is the same as the depth of penetration of the needle. A needle outlet that is either formed by a bevel or by an opening through the side of the needle has a measurable exposed height. It is understood that a single needle may have more than one opening or outlets suitable for delivery of substances to the dermal space.

[0047] It has also been found that by controlling the pressure of injection or infusion may avoid the high backpressure exerted during ID administration. By placing a constant pressure directly on the liquid interface a more constant delivery rate can be achieved, which may optimize absorption and obtain the improved pharmacokinetics. Delivery rate and volume can also be controlled to prevent the formation of wheals at the site of delivery and to prevent backpressure from pushing the dermal-access means out of the skin. The appropriate delivery rates and volumes

to obtain these effects for a selected substance may be determined experimentally using only ordinary skill. Increased spacing between multiple needles allows broader fluid distribution and increased rates of delivery or larger fluid volumes. In addition, it has been found that ID infusion or injection often produces higher initial plasma levels of drug than conventional SC administration, particularly for drugs that are susceptible to in vivo degradation or clearance or for compounds that have an affinity to the SC adipose tissue or for macromolecules that diffuse slowly through the SC matrix. This may, in many cases, allow for smaller doses of the substance to be administered via the ID route.

[0048] The administration methods useful for carrying out the invention include both bolus and infusion delivery of drugs and other substances to humans or animals subjects. A bolus dose is a single dose delivered in a single volume unit over a relatively brief period of time, typically less than about 10 minutes. Infusion administration comprises administering a fluid at a selected rate that may be constant or variable, over a relatively more extended time period, typically greater than about 10 minutes. To deliver a substance the dermal-access means is placed adjacent to the skin of a subject providing directly targeted access within the intradermal space and the substance or substances are delivered or administered into the intradermal space where they can act locally or be absorbed by the bloodstream and be distributed systematically. The dermal-access means may be connected to a reservoir containing the substance or substances to be delivered. The form of the substance or substances to be delivered or administered include solutions thereof in pharmaceutically acceptable diluents or solvents, emulsions, suspensions, gels, particulates such as micro- and nanoparticles either suspended or dispersed, as well as in-situ forming vehicles of the same. Delivery from the reservoir into the intradermal space may occur either passively, without application of the external pressure or other driving means to the substance or substances to be delivered, and/or actively, with the application of pressure or other driving means. Examples of preferred pressure generating means include pumps, syringes, elastomer membranes, gas pressure, piezoelectric, electromotive, electromagnetic pumping, or Belleville springs or

washers or combinations thereof. If desired, the rate of delivery of the substance may be variably controlled by the pressure-generating means. As a result, the substance enters the intradermal space and is absorbed in an amount and at a rate sufficient to produce a clinically efficacious result.

[0049] As used herein, the term "clinically efficacious result" is meant a clinically useful biological response including both diagnostically and therapeutically useful responses, resulting from administration of a substance or substances. For example, diagnostic testing or prevention or treatment of a disease or condition is a clinically efficacious result. Such clinically efficacious results include diagnostic results such as the measurement of glomerular filtration pressure following injection of inulin, the diagnosis of adrenocortical function in children following injection of ACTH, the causing of the gallbladder to contract and evacuate bile upon injection of cholecystokinin and the like as well as therapeutic results, such as clinically adequate control of blood sugar levels upon injection of insulin, clinically adequate management of hormone deficiency following hormone injection such as parathyroid hormone or growth hormone, clinically adequate treatment of toxicity upon injection of an antitoxin and the like .

[0050] Substances that can be delivered intradermally in accordance with the present invention are intended to include pharmaceutically or biologically active substances including include diagnostic agents, drugs, and other substances which provide therapeutic or health benefits such as for example nutraceuticals. Diagnostic substances useful with the present invention include macromolecular substances such as, for example, inulin, ACTH (e.g. corticotropin injection), luteinizing hormone-releasing hormone (eg., Gonadorelin Hydrochloride), growth hormone-releasing hormone (e.g. Sermorelin Acetate), cholecystokinin (Sincalide), parathyroid hormone and fragments thereof (e.g. Teriparatide Acetate), thyroid releasing hormone and analogs thereof (e.g. protirelin), secretin and the like.

[0051] Therapeutic substances which can be used with the present invention include Alpha-1 anti-trypsin, Anti-Angiogenesis agents, Antisense, butorphanol, Calcitonin and analogs, Ceredase, COX-II inhibitors, dermatological agents, dihydroergotamine, Dopamine agonists and antagonists, Enkephalins and other opioid peptides, Epidermal growth factors, Erythropoietin and analogs, Follicle stimulating hormone, G-CSF, Glucagon, GM-CSF, granisetron, Growth hormone and analogs (including growth hormone releasing hormone), Growth hormone antagonists, Hirudin and Hirudin analogs such as Hirulog, IgE suppressors, Insulin, insulinotropin and analogs, Insulin-like growth factors, Interferons, Interleukins, Luteinizing hormone, Luteinizing hormone releasing hormone and analogs, Heparins, Low molecular weight heparins and other natural, modified, or synthetic glycoaminoglycans, M-CSF, metoclopramide, Midazolam, Monoclonal antibodies, Peglyated antibodies, Pegylated proteins or any proteins modified with hydrophilic or hydrophobic polymers or additional functional groups, Fusion proteins, Single chain antibody fragments or the same with any combination of attached proteins, macromolecules, or additional functional groups thereof, Narcotic analgesics, nicotine, Non-steroid anti-inflammatory agents, Oligosaccharides, ondansetron, Parathyroid hormone and analogs, Parathyroid hormone antagonists, Prostaglandin antagonists, Prostaglandins, Recombinant soluble receptors, scopolamine, Serotonin agonists and antagonists, Sildenafil, Terbutaline, Thrombolytics, Tissue plasminogen activators, TNF - , and TNF - antagonist, the vaccines, with or without carriers/adjuvants, including prophylactics and therapeutic antigens (including but not limited to subunit protein, peptide and polysaccharide, polysaccharide conjugates, toxoids, genetic based vaccines, live attenuated, reassortant, inactivated, whole cells, viral and bacterial vectors) in connection with, addiction, arthritis, cholera, cocaine addiction, diphtheria, tetanus, HIB, Lyme disease, meningococcus, measles, mumps, rubella, varicella, yellow fever, Respiratory syncytial virus, tick borne Japanese encephalitis, pneumococcus, streptococcus, typhoid, influenza, hepatitis, including hepatitis A, B, C and E, otitis media, rabies, polio, HIV, parainfluenza, rotavirus, Epstein Barr Virus, CMV, chlamydia, non-typeable haemophilus, moraxella catarrhalis, human papilloma virus, tuberculosis including BCG, gonorrhoea, asthma,

atherosclerosis malaria, E-coli, Alzheimer's Disease, H. Pylori, salmonella, diabetes, cancer, herpes simplex, human papilloma and the like other substances including all of the major therapeutics such as agents for the common cold, Anti-addiction, anti-allergy, anti-emetics, anti-obesity, antiosteoporotic, anti-infectives, analgesics, anesthetics, anorexics, antiarthritics, antiasthmatic agents, anticonvulsants, antidepressants, antidiabetic agents, antihistamines, anti-inflammatory agents, antimigraine preparations, anti-motion sickness preparations, anti-nauseants, antineoplastics, antiparkinsonism drugs, antipruritics, antipsychotics, antipyretics, anticholinergics, benzodiazepine antagonists, vasodilators, including general, coronary, peripheral and cerebral, bone stimulating agents, central nervous system stimulants, hormones, hypnotics, immunosuppressives, muscle relaxants, parasympatholytics, parasympathomimetics, prostaglandins, proteins, peptides, polypeptides and other macromolecules, psychostimulants, sedatives, and sexual hypofunction and tranquilizers.

[0052] Pharmacokinetic analysis of insulin infusion data was carried out as follows. Stepwise nonlinear least-squares regression was used to analyze the insulin concentration-time data from each individual animal. Initially, an empirical biexponential equation was fit to the insulin concentration-time data for the negative control condition. This analysis assumed first-order release of residual insulin, and recovered parameters for the first-order rate constant for release, the residual insulin concentration at the release site, a lag time for release, and a first-order rate constant for elimination of insulin from the systemic circulation. The parameters recovered in this phase of the analysis are of no intrinsic importance, but merely account for the fraction of circulating insulin derived from endogenous sources.

[0053] The second step of the analysis involved fitting an explicit compartmental model to the insulin concentration-time data during and after subcutaneous or intradermal infusion. The scheme upon which the mathematical model was based is shown in the upper part of Figure 1.[PK/PD model fig]. Infusion of insulin proceeded from $t = 0$ to $t = 240$ min; after a lag time ($t_{lag,2}$), absorption from the infusion site was

mediated by a first-order process governed by the absorption rate constant k_a . Insulin absorbed into the systemic circulation distributed into an apparent volume V contaminated by an unknown fractional bioavailability F , and was eliminated according to a first-order rate constant K . The fitting routine recovered estimates of $t_{lag,2}$, k_a , V/F , and K ; parameters associated with the disposition of endogenous insulin (C_R , $t_{lag,1}$, k_R), which were recovered in the first step of the analysis, were treated as constants.

[0054] Parameter estimates are reported as mean \pm SD. The significance of differences in specific parameters between the two different modes of insulin administration (subcutaneous versus intradermal infusion) was assessed with the paired Student's t -test.

Pharmacodynamic analysis of insulin infusion data was calculated as follows. Plasma concentrations of glucose were used as a surrogate for the pharmacologic effect of insulin. The change in response variable R (plasma glucose concentration) with respect to time t was modeled as

$$\frac{dR}{dt} = k_{in} - E \cdot k_{out}$$

[0055] where k_{in} is the zero-order infusion of glucose, k_{out} is the first-order rate constant mediating glucose elimination, and E is the effect of insulin according to the sigmoidal Hill relationship

$$E = \frac{E_{max} \cdot C^\gamma}{EC_{50}^\gamma + C^\gamma}$$

[0056] in which E_{max} is the maximal stimulation of k_{out} by insulin, EC_{50} is the insulin concentration at which stimulation of k_{out} is half maximal, C is the concentration of insulin, and γ is the Hill coefficient of the relationship. Initial modeling efforts utilized the plasma concentration of insulin as the mediator of

pharmacologic response. However, this approach did not capture the delay in response of plasma glucose to increasing concentrations of plasma insulin. Therefore, an effect-compartment modeling approach was finally adopted in which the effect of insulin was mediated from a hypothetical effect compartment peripheral to the systemic pharmacokinetic compartment

[0057] The pharmacodynamic analysis was conducted in two steps. In the first step of the analysis, initial estimates of the pharmacokinetic parameters associated with the disposition of glucose (k_{out} and the volume of distribution of glucose, $V_{glucose}$) were determined from the glucose concentration-time data in the negative control condition. The full integrated pharmacokinetic-pharmacodynamic model then was fit simultaneously to the glucose concentration-time data from the negative control condition and each insulin delivery condition for each animal (i.e., two sets of pharmacodynamic parameters were obtained for each animal: one from the simultaneous analysis of the subcutaneous insulin infusion/negative control data, and one from the simultaneous analysis of the intradermal insulin infusion/negative control data). In all pharmacodynamic analyses, the parameters governing insulin disposition obtained during pharmacokinetic analysis of insulin concentration-time data from each animal were held constant.

[0058] All other pharmacokinetic analyses were calculated using non-compartmental methods using similar software programs and techniques known in the art..

[0059] Having described the invention in general, the following specific but not limiting examples and reference to the accompanying Figures set forth various examples for practicing the dermal accessing, direct targeting drug administration method and examples of dermally administered drug substances providing improved PK and PD effects.

[0060] A representative example of dermal-access microdevice comprising a single needle were prepared from 34 gauge steel stock (MicroGroup, Inc., Medway, MA) and a single 28° bevel was ground using an 800 grit carborundum grinding wheel. Needles were cleaned by sequential sonication in acetone and distilled water, and flow-checked with distilled water. Microneedles were secured into small gauge catheter tubing (Maersk Medical) using UV-cured epoxy resin. Needle length was set using a mechanical indexing plate, with the hub of the catheter tubing acting as a depth-limiting control and was confirmed by optical microscopy. For experiments using needles of various lengths, the exposed needle lengths were adjusted to 0.5, 0.8, 1, 2 or 3 mm using the indexing plate. Connection to the fluid metering device, either pump or syringe, was via an integral Luer adapter at the catheter inlet. During injection, needles were inserted perpendicular to the skin surface, and were either held in place by gentle hand pressure for bolus delivery or held upright by medical adhesive tape for longer infusions. Devices were checked for function and fluid flow both immediately prior to and post injection. This Luer Lok single needle catheter design is hereafter designated SS1_34.

[0061] Yet another dermal-access array microdevices was prepared consisting of 1" diameter disks machined from acrylic polymer, with a low volume fluid path branching to each individual needle from a central inlet. Fluid input was via a low volume catheter line connected to a Hamilton microsyringe, and delivery rate was controlled via a syringe pump. Needles were arranged in the disk with a circular pattern of 15 mm diameter. Three-needle and six-needle arrays were constructed, with 12 and 7 mm needle-to-needle spacing, respectively. All array designs used single-bevel, 34 G stainless steel microneedles of 1 mm length. The 3-needle 12mm spacing catheter-design is hereafter designated SS3_34B, 6-needle 7mm spacing catheter-design is hereafter designated SS6_34A.

[0062] Yet another dermal-access array microdevices was prepared consisting of 11mm diameter disks machined from acrylic polymer, with a low volume fluid path branching to each individual needle from a central inlet. Fluid input was via a low

volume catheter line connected to a Hamilton microsyringe, and delivery rate was controlled via a syringe pump. Needles were arranged in the disk with a circular pattern of about 5 mm diameter. Three-needle arrays of about 4 mm spacing connected to a catheter as described above. These designs are hereafter designated SS3S_34_1, SS3C_34_2, and SS3S_34_3 for 1mm, 2mm, and 3mm needle lengths respectively.

[0063] Yet another dermal-access ID infusion device was constructed using a stainless steel 30 gauge needle bent at near the tip at a 90-degree angle such that the available length for skin penetration was 1-2 mm. The needle outlet (the tip of the needle) was at a depth of 1.7-2.0 mm in the skin when the needle was inserted and the total exposed height of the needle outlet 1.0-1.2 mm. This design is hereafter designated SSB1_30.

[0064] EXAMPLE I

[0065] Slow-infusion ID insulin delivery was demonstrated in swine using a hollow, silicon-based single-lumen microneedle (2 mm total length and 200 X 100 μ m OD, corresponding to about 33 gauge) with an outlet 1.0 μ m from the tip (100 μ m exposed height), was fabricated using processes known in the art (US Patent No. 5,928,207) and mated to a microbore catheter (Disetronic). The distal end of the microneedle was placed into the plastic catheter and cemented in place with epoxy resin to form a depth-limiting hub. The needle outlet was positioned approximately 1 mm beyond the epoxy hub, thus limiting penetration of the needle outlet into the skin to approximately 1 mm., which corresponds to the depth of the intradermal space in swine.. The catheter was attached to a MiniMed 507 insulin pump for control of fluid delivery. The distal end of the microneedle was placed into the plastic catheter and cemented in place with epoxy resin to form a depth-limiting hub. The needle outlet was positioned approximately 1 mm beyond the epoxy hub, thus limiting penetration of the needle outlet into the skin to approximately 1 mm., which corresponds to the depth of the intradermal space in swine. The patency of the fluid flow path was confirmed by visual observation, and no obstructions were observed at pressures

generated by a standard 1-cc syringe. The catheter was connected to an external insulin infusion pump (MiniMed 507) via the integral Luer connection at the catheter outlet. The pump was filled with Humalog™ (Lispro) insulin (Eli Lilly, Indianapolis, IN) and the catheter and microneedle were primed with insulin according to the manufacturer's instructions. Sandostatin® (Sandoz, East Hanover, NJ) solution was administered via IV infusion to anesthetized swine to suppress basal pancreatic function and insulin secretion. After a suitable induction period and baseline sampling, the primed microneedle was inserted perpendicular to the skin surface in the flank of the animal up to the hub stop. Insulin infusion at a rate of 2 U/hr was used and maintained for 4 hr. Blood samples were periodically withdrawn and analyzed for serum insulin concentration and blood glucose values. Baseline insulin levels before infusion were at the background detection level of the assay. After initiation of the infusion, serum insulin levels showed an increase that was commensurate with the programmed infusion rates. Blood glucose levels also showed a corresponding drop relative to negative controls (NC) without insulin infusion and this drop was improved relative to conventional SC infusion. In this experiment, the microneedle was demonstrated to adequately breach the skin barrier and deliver a drug *in vivo* at pharmaceutically relevant rates. The ID infusion of insulin was demonstrated to be a pharmacokinetically acceptable administration route, and the pharmacodynamic response of blood glucose reduction was also demonstrated. Calculated PK parameters for ID infusion indicate that insulin is absorbed much faster than via than SC administration. Absorption from the ID space begins almost immediately: the lag time prior to absorption (t_{lag}) was 0.88 vs. 13.6 min for ID and SC respectively. Also the rate of uptake from the administration site is increased by approximately 3-fold, $k_a = 0.0666$ vs. 0.0225 min^{-1} for ID and SC respectively. The bioavailability of insulin delivered by ID administration is increased approximately 1.3 fold greater than SC administration.

[0066] EXAMPLE II

[0067] Bolus delivery of Lilly Lispro fast acting insulin was performed using ID and SC bolus administration. The ID injection microdevice was dermal access array

design SS3_34. 10 international insulin units (U) corresponding to 100 uL volume respectively, were administered to diabetic Yucatan Mini swine. Test animals had been previously been rendered diabetic by chemical ablation of pancreatic islet cells, and were no longer able to secrete insulin. Test animals received their insulin injection either via the microneedle array or via a standard 30 G X ½ in. needle inserted laterally into the SC tissue space. Circulating serum insulin levels were detected using a commercial chemiluminescent assay kit (Immulite, Los Angeles, CA) and blood glucose values were determined using blood glucose strips. ID injections were accomplished via hand pressure using an analytical microsyringe and were administered over approximately 60 sec. By comparison, SC dosing required only 2-3 sec. Referring to Figure 1, it is shown that serum insulin levels after bolus administration demonstrate more rapid uptake and distribution of the injected insulin when administered via the ID route. The time to maximum concentration (T_{max}) is shorter and the maximum concentration obtained (C_{max}) is higher for ID vs. SC administration. In addition, Figure 2 also demonstrates the pharmacodynamic biological response to the administered insulin, as measured by the decrease in blood glucose (BG), showed faster and greater changes in BG since more insulin was available early after ID administration.

[0068] EXAMPLE III

[0069] Lilly Lispro is regarded as fast acting insulin, and has a slightly altered protein structure relative to native human insulin.. Hoechst regular insulin, maintains the native human insulin protein structure that is chemically similar, but has slower uptake than Lispro when administered by the traditional SC route. Both insulin types were administered in bolus via the ID route to determine if any differences in uptake would be discernable by this route. 5U of either insulin type were administered to the ID space using dermal access microdevice design SS3_34. The insulin concentration verses time data shown in Figure 3. When administered by the ID route the PK profiles for regular and fast-acting insulin were essentially identical, and both insulin types exhibited faster uptake than Lispro given by the traditional SC route. This is evidence that the uptake mechanism for ID administration is less affected by minor

biochemical changes in the administered substance, and that ID delivery provides an advantageous PK uptake profile for regular insulin that is superior to SC administered fast-acting insulin.

[0070] EXAMPLE IV

[0071] Bolus delivery of Lilly Lispro fast-acting insulin via microneedle arrays having needles of various lengths was conducted to demonstrate that the precise deposition of drug into the dermal space is necessary to obtain the PK advantages and distinctions relative to SC. Thus, 5U of Lilly Lispro fast-acting insulin was administered using dermal access design SS3_34. Additional microdevices of the same needle array configuration were fabricated whereby exposed needle lengths of the microdevice array were lengthened to include arrays with needles lengths of 2 and 3 mm. The average total dermal thickness in Yucatan Mini swine ranges from 1.5-2.5 mm. Therefore insulin deposition is expected to be into the dermis, approximately at the dermal/SC interface, and below the dermis and within the SC for 1mm, 2mm, and 3mm length needles respectively. Bolus insulin administration was as described in EXAMPLE II.. Average insulin concentrations verses time are shown in Figure 4. The data clearly shows as microneedle length is increased, the resulting PK profile begins to more closely resemble SC administration. This data demonstrates the benefits of directly targeting the dermal space, such benefits include rapid uptake and distribution, and high initial concentrations. Since the data are averages of multiple examples, they do not show the increased inter-individual variability in PK profiles from longer 2 and 3mm microneedles. This data demonstrates that since skin thickness may vary both between individuals and even within a single individual, shorter needle lengths that accurately target the dermal space are more reproducible in their PK profile since they are depositing the drug more consistently in the same tissue compartment. This data demonstrates longer microneedles that deposit or administer substances deeper into the dermal space, or partially or wholly into the SC space, mitigate or eliminate the PK advantages in comparison to shallow, directly targeted administrations to the highly vascularized dermal region.

[0072] EXAMPLE V

[0073] Bolus delivery of Lantus long-acting insulin was delivered via the ID route . Lantus is an insulin solution that forms microprecipitates at the administration site upon injection. These microparticulates undergo slow dissolution within the body to provide (according to the manufacturer's literature) a more stable low level of circulating insulin than other current long-acting insulin such as crystalline zinc precipitates (e.g. Lente, NPH). Lantus insulin (10 U dose, 100 uL) was administered to diabetic Yucatan Mini pigs using the dermal access design SS3_34 and by the standard SC method as previously described. Referring to Figure 5, when administered via the ID route, similar PK profiles were obtained relative to SC. Minor distinctions include a slightly higher "burst" immediately after the ID insulin delivery. This demonstrates that the uptake of even very high molecular weight compounds or small particles is achievable via ID administration. More importantly this supports the fact that the biological clearance mechanism in the body is not appreciably changed by the administration route, nor is the way in which that the drug substance is utilized. This is extremely important for drugs compounds that have a long circulating half-life (examples would be large soluble receptor compounds or other antibodies for cancer treatment, or chemically modified species such as PEGylated drugs). NEED CLAIM

[0074] EXAMPLE VI

[0075] Bolus ID delivery of human granulocyte colony stimulating factor (GCSF) (Neupogen) was administered via dermal access microdevice designs SS3_34B (array) or SS1_34 (single needle) to Yucatan minipigs. Delivery rate was controlled via a Harvard syringe pump and was administered over a 1-2.5 min period. Figure 6 shows the PK availability of GCSF in blood plasma as detected by an ELISA immunoassay specific for GCSF. Administration via IV and SC delivery was performed as controls. Referring to Figure 6 bolus ID delivery of GCSF shows the more rapid uptake associated with ID delivery. C_{max} is achieved at approximately 30-90 minutes vs. 120 min for SC. Also the bioavailability is dramatically increased by

an approximate factor of 2 as evidenced by the much higher area under the curve (AUC). Circulating levels of GCSF are detectable for an extended period, indicating that ID delivery does not alter the intrinsic biological clearance mechanism or rate for the drug. These data also show that device design has minimal effect on the rapid uptake of drug from the ID space. The data referred to in Figure 7 also shows the degree and time course of white blood cell expansion as a result of GCSF administration with respect to a negative control (no GCSF administered). White blood cell (WBC) counts were determined by standard cytometric clinical veterinary methods. ID delivery exhibits the same clinically significant biological outcomes. Although all delivery means give approximately equal PD outcomes, this data suggests ID delivery could enable use half the dose to achieve essentially the same physiological result in comparison to SC, due to approximately 2-fold bioavailability increase.

[0076] EXAMPLE VII

An ID administration experiment was conducted using a peptide drug entity: human parathyroid hormone 1-34 (PTH). PTH was infused for a 4 h period, followed by a 2 h clearance. Control SC infusion was through a standard 31-gauge needle inserted into the SC space lateral to the skin using a "pinch-up" technique. ID infusion was through dermal access microdevice design SSB1_30 (a stainless steel 30-gauge needle bent at the tip at a 90° angle such that the available length for skin penetration was 1-2 mm). The needle outlet (the tip of the needle) was at a depth of 1.7-2.0 mm in the skin when the needle was inserted. A 0.64 mg/mL PTH solution was infused at a rate of 75 μ L/hr. Flow rate was controlled via a Harvard syringe pump. Weight normalized PTH plasma levels are shown in Figure XX. The weight normalized delivery profiles show a larger area under the curve (AUC) indicating higher bioavailability, higher peak values at earlier sampling timepoints (e.g. 15 and 30 min) indicating more rapid onset from ID delivery, and rapid decrease following termination of infusion (also indicative of rapid uptake without a depot effect).

Table 2. Calculated LMWHPK Data

Condition:	SC		1.0 mm microneedle		0.5 mm microneedle	
	Mean	SD	Mean	SD	Mean	SD
t_{\max} (h)	3.0	3.6	1.0	0.3	0.8	0.3
C_{\max} (IU/mL)	0.6	0.3	1.1	0.1	1.5	0.3

[0077] EXAMPLE VIII

[0078] Referring to Figure 8, representative weight normalized plasma profiles following bolus delivery of Fragmin, low molecular weight heparin fragment(LMWH) in Yucatan mini-pigs via various dermal access microdevice configurations are presented. In each case the delivered dose was 2500 IU (international units) of Fragmin (100 ul of a 25000 IU/mL formulation). Standard SC delivery was performed via a standard 30 G needle inserted laterally into the SC tissue space via a pinch-up technique. Dermal access microdevice designs SS1_34 of 0.5 or 1.0mm needle length connected to catheter tubing were used for dosing. During use the fully exposed length of microneedle was inserted perpendicularly to the skin surface up to the depth-limiter and held in place by mechanical means for the duration of drug instillation. The microneedle bolus injection was via hand pressure from a glass microsyringe over a 1-2.5 min period. The calculated pharmacokinetic results of Table 1 show the increased C_{\max} and decreased T_{\max} resulting from microdevice delivery. The profiles obtained from both microneedle devices was essentially equivalent indicating that the delivery profile is essentially independent of device configuration providing the device appropriately accesses and delivers the drug substance within the dermal tissue compartment. Equivalent changes in pharmacokinetic uptake can be generated using the other dermal access microdevice systems including arrays composed of 3 and 6 microneedles with the same dimensions and seating depths indicated above.

[0079] EXAMPLE IX

[0080] Referring to Figure 9 showing comparative plasma profiles for bolus administered Fragmin dosing conditions 1). SC 100uL injected volume; 2500IU total dose, 2). ID 100uL injected volume; 2500IU total dose; 1.0mm needle length (SS1_34) and 3). ID 100uL injected volume; 2500IU total dose; 0.5mm needle length (SS1_34). At the time of dosing, these animals were weight-matched, within a 8.8 to 12.3 kg weight range. All plasma profiles have been normalized to an average animal weight of 15.0 kg, by multiplying the raw data by the animal weight at time of dosing and dividing by 15. However, individual plasma profiles are not adjusted for dosing variability. PK parameters are calculated based on the raw data, and are corrected both for dosing levels and animal weight. This data demonstrates the reduced onset time for drug bioavailability and distribution for ID administration compared to SC.

[0081] EXAMPLE X

[0082] Referring to Figure 9 which shows representative weight normalized plasma profiles of short infusion delivery of Fragmin LMWH in Yucatan mini-pigs. A total of 2500 IU in a 200 uL volume (12500 IU/mL concentration) of LMWH was infused over durations ranging from 0.5-2.0 h. The volumetric infusion rate ranged between 100-400 uL/h. The dermal access array microdevice was of design SS3_34 connected to a syringe pump for control of fluid delivery. Each microneedle in the array had a 1 mm extended length for insertion. ID bolus injection of an equivalent dose (100 uL of 25000 IU/ml) LMWH over a <2 min period via a similar microneedle array and standard SC bolus administration are shown for comparison. The resulting plasma profiles demonstrate the highly controllable drug delivery profiles obtainable with a microdevice intradermal system. This data demonstrates the infusion control means allows for modulation of the pharmacokinetics via the infusion rate. As volumetric infusion rates decrease, C_{max} and T_{max} decrease and increase, respectively. Within experimental error T_{max} for Fragmin was routinely obtained at the cessation of the infusion period. This short infusion administration result demonstrates the ability to deliver greater than normal total fluid volumes than standard ID administrations (Mantoux technique is limited to about 100 to 150uL/dose)..

[0083] EXAMPLE XI

[0084] Referring to Figure 10, which shows representative weight normalized plasma profiles following slow infusion delivery of Fragmin LMWH in Yucatan mini-pigs. A total of 2000 IU in an 80 uL volume (25000 IU/mL concentration) of LMWH was infused over a 5 hour period. The volumetric infusion rate was 16 uL/h. The infusion means was a commercial insulin pump connected to either an ID microdevice of design SS1_34, or a commercial insulin infusion catheter. The resulting plasma profiles again indicate the more rapid onset of LMWH infused via microdevices. After removal of the catheter set at 5 hours, the ID delivery also exhibits the lack of depot effect, as evidenced by the immediate decline of detectable plasma activity. In contrast, the plasma levels of SC infused LMWH do not peak until 7 h, fully 2 h after infusion cessation. Neither infusion method reaches steady state over the experimental duration, but this was previously predicted via PK modeling. This example readily demonstrates that the PK advantages of controlled ID delivery are available at low infusion rates, and the degree of control, which can be achieved in dosing profiles. This particular profile would be optimal for drugs such as LMWH, insulin, etc that require low continuous circulating basal levels without high peak concentrations.

[0085] EXAMPLE XII

[0086] Referring to Table 2 which shows weight normalized serum levels of hGH after bolus delivery of Genotropin recombinant human growth hormone via intradermal microdevices and standard subcutaneous injection methods of 3.6 IU of Genotropin. Injection volume was 100 uL and the drug concentration was 36 IU/mL. Dermal access array microdevices were SS1_34 and SS3_34 designs with 1mm exposed needle length. The rate of microdevice injection for both single and three-needle arrays was controlled at 45 uL/min using a syringe pump, for a nominal bolus infusion duration of 2.22 minutes. SC delivery was via a 27 G insulin catheter, at a 1.0 mL/min flow rate, for a nominal 10 sec injection. The resultant pharmacokinetic distinctions are clearly evident, with ID delivery resulting in drastically decreased

t_{max} , and much increased C_{max} . Biological half life, and bioavailability are statistically equivalent for both ID and SC routes.. Administration by either single needle or array intradermal dermal access microdevice configurations produce equivalent pharmacokinetic performance.

Table 2: Calculated PK parameters for Genotropin administration

PK parameters	SC	ID single needle	ID 6-needle array
<i>Dose (IU/kg)</i>	<i>0.161</i> <i>±0.01</i>	<i>0.164±0.</i> <i>01</i>	<i>0.160±0.0</i> <i>2</i>
<i>C_{max} (mIU/L)</i>	<i>158.5</i> <i>±31.0</i>	<i>612.6±1</i> <i>87.1</i>	<i>582.1±39</i> <i>1.0</i>
<i>t_{max} (h)</i>	<i>2.75±</i> <i>0.46</i>	<i>0.47±0.2</i> <i>5</i>	<i>0.63±0.23</i>
<i>t_{1/2z} (h)</i>	<i>1.19±</i> <i>0.49</i>	<i>2.02±0.4</i> <i>8</i>	<i>1.71±0.43</i>
<i>AUC_{INF(pred)}</i> <i>(mIU x h/L)</i>	<i>920.2</i> <i>±251.</i> <i>7</i>	<i>850.0±1</i> <i>70.0</i>	<i>847.4±33</i> <i>2.3</i>
<i>F (%)</i>	<i>114.6</i>	<i>104.0</i>	<i>101.7</i>

[0087] EXAMPLE XIII

[0088] Referring to the data in Table 3, bolus delivery of Almotriptan, a low molecular weight, highly water soluble antimigraine compound, via intradermal microdevices and standard subcutaneous methods demonstrated statistically equivalent PK profiles. The table below shows calculated PK parameters determined from measured serum levels after injection of 3.0 mg of almotriptan. Injection volume for both SC and ID was 100 uL and the drug concentration was 30 mg/mL. Microdevices designs SS3_34 and SS6_34 were used administered over about 2-2.5

minutes. Almotriptan is a small hydrophilic compound that shows no apparent depot from SC injection. Therefore, differences in the pharmacokinetic uptake between ID and SC administration were not observed. This drug substance can readily partition through the tissue space for rapid absorption via either route. However, ID administration may still be adventitious for reduced patient perception and ready and rapid access to an appropriate administration site.

[0089] Table 3: Mean (\pm standard deviation) almotriptan PK parameters following SC and ID administration

Parameters	SC	ID (single)	ID (array)
AUC _{0-∞} (ng h/mL)	55.9 (6.04)	53.3 (15.7)	54.6 (14.0)
Clearance (L/hr)	55.1 (5.87)	60.1 (15.3)	58.7 (12.7)
C _{max} (ng/mL)	61.0 (19.4)	63.6 (26.1)	77.2 (54.2)
t _{max} (h)	0.13 (0.05)	0.14 (0.08)	0.16 (0.08)
z (h ⁻¹)	0.36 (0.04)	0.36 (0.08)	0.31 (0.08)
t _{1/2} (h)	1.95 (0.23)	2.03 (0.46)	2.39 (0.64)

[0090] The above examples and results demonstrate the inventive delivery method using multi-point array ID administration and single needle ID administration results in more rapid uptake with higher C_{max} than SC injection. ID uptake and distribution is ostensibly unaffected by device geometry parameters, using needle lengths of about 0.5 to about 1.7mm, needle number and needle spacing. No concentration limit for biological absorption was found and PK profiles were dictated principally by the concentration-based delivery rate. The primary limitations of ID administration are the total volume and volumetric infusion-rate limits for leak-free instillation of exogenous substances into a dense tissue compartment. Since absorption of drugs from the ID space appears to be insensitive to both device design and volumetric infusion rate, numerous formulation/device combinations can be used to overcome this limitations and provide the required or desired therapeutic profiles.

For example, volume limited dosing regimens can be circumvented either by using more concentrated formulations or increasing the total number of instillation sites. In addition, effective PK control is obtained by manipulating infusion or administration rate of substances.

[0091] In general, ID delivery as taught by the methods described hereto via dermal access microneedle devices provides a readily accessible and reproducible parenteral delivery route, with high bioavailability, as well as the ability to modulate plasma profiles by adjusting the device infusion parameters, since uptake is not rate-limited by biological uptake parameters.

[0092] In the previously described examples, the methods practiced by the invention demonstrate the ability to deliver a drug in vivo with greatly improved pharmaceutically relevant rates. This data indicates an improved pharmacological result for ID administration as taught by the methods described of other drugs in humans would be expected according to the methods of the invention.